

RETAINED PRIMITIVE REFLEX SCREENING

Name: _____ DOB: _____ Date

REFLEX & COMMON SIGNS OF ABNORMALITY	NAD	Trace	Mild	Moderate	Significant	Notes
Fear Paralysis Reflex						
inability to make or make or maintain eye contact						
stares without blinking						
holds breath						
tension up whole body						
clenches fists						
sways						
toes curl						
Babkin Response						
fingers close						
withdrawing the arm						
tension in trunk						
mouth moves or opens						
head turns to same side as stimulated palm						
tongue moves						
fingers on other hand move						
Moro (vestibular)						
<input type="checkbox"/> Supine <input type="checkbox"/> Standing						
movement of arms outward						
dorsiflexion of toes						
distress, does not like test						
withdrawal						
quick inhale						
change in face color						
chin tucks forward						
loss of tone in core						
Palmar						
wiggles fingers or thumb						
moves fingers or thumb inward						
ticklish, irritating or painful						
thumb and fingers close or extend on stimulation						

RETAINED PRIMITIVE REFLEX SCREENING

Name: _____ DOB: _____ Date _____

Tonic Labyrinthine Reflex (TLR)-forward						
loss of balance						
compensatory change in muscle tone at knees						
instability of upper body						
instability of lower body						
dizziness, nausea						
swaying						
TLR-backward						
loss of Balance						
compensatory change in muscle tone at knees						
instability of upper body						
instability of lower body						
dizziness, nausea						
swaying						
TLR-Prone						
flexion of lower extremities						
flexion of upper extremities						
difficulty lifting head or arms						
difficulty lifting legs						
inability to hold position for 3-5 seconds						
holds breath						
Asymmetrical Tonic Neck Reflex (ATNR)--Schilder's test						
movement in the body or arms on the side to which the head is turned						
tendency to increased extensor tone on the side to which the head is turned						
inability to relax the neck muscles						
ATNR-Ayers						
elbow bends on occipital side						
elbow extends on face side						
shoulder drops						
instability in trunk or hips						
feet raise, cross, or kick						

RETAINED PRIMITIVE REFLEX SCREENING

Name: _____ DOB: _____ Date _____

Spinal Galant Reflex						
movement of hip outwards with stimulation						
lifting of a foot						
drop or elevation of shoulder						
flexion or extension of supporting arm						
Rooting Reflex						
twitching of mouth in response to stimulus						
withdrawal						
mouth opens						
hands move						
wipes mouth after test						
turn head towards stimulus						
Babinski Reflex						
walks on outside of feet						
flat footed -feet flop outwards						
feet point to each other						
feet point rigidly upward						
toes fan out with stimulation						
withdraw with stimulus						
Labyrinthine Head Righting Reflex (LHRR)						
head follows displacement of body						
head overcompensates on retrieval to midline						
Oculo-Head Righting Reflex (OHRR)						
head follows displacement of body						
head overcompensates on retrieval to midline						
STNR						
tremor in one or both arms						
instability in trunk or hips						
bending of arms as a result of head flexion and/or raising of feet						
straightening of arms and flexion of knees as a result of head extension						
sway or arching of back						
sitting on feet						